



# Specimen Label

## PRINCIPAL FUNCTIONING AGENTS:

A proprietary blend of aliphatic hydrocarbons, fatty acids and alkyl ethoxylates.....100%

**Total** ..... **100%**

All ingredients are exempt from tolerance under 40 CFR 180.

## KEEP OUT OF REACH OF CHILDREN CAUTION

Read label before use. Keep container tightly closed. Keep only in original container.

Do not eat, drink or smoke when using this product. Avoid breathing dust, fume, gas, mist, vapors or spray. Wash hands thoroughly after handling. Wear protective gloves, protective clothing, eye protection and face protection.

## GENERAL INFORMATION

**ELITE BARREN** is formulated with a superior surfactant system and paraffinic oil to improve deposition and penetration of the active ingredient or nutrients into the target surface.

**ELITE BARREN** promotes efficacy by increasing soil adsorption which can help to reduce run-off and leaching of the spray application.

**ELITE BARREN** is recommended for application in a broad range of crop and non-crop uses including agriculture, turf and ornamental, industrial vegetation management, and specialty crops.

May be used with herbicides, fungicides, or insecticides.

## DIRECTIONS FOR USE

Follow the instructions on the most restrictive active ingredient label.

Tank mixtures should only be applied within the label recommendations of every product in the tank mixture.

Caution should be used when applying this or any other product on crops during hot, dry conditions.

Though phytotoxicity has been evaluated, care should be taken by conducting a jar test or testing this product on sensitive plants before full application is performed.

## Order of addition

Prior to mixing, fill the spray tank with half of the carrier. Start the sprayer and check to make sure that all valves and gauges work and that there is adequate tank agitation.

In general, follow the W-A-L-E-S plan when adding herbicides to a tank mix.

1. Wettable Powders (WP) then Flowables (F, DF)
2. Agitate then add anti-foaming compounds or buffers.
3. Liquid and Soluble products
4. Emulsifiable concentrates (EC)
5. **ELITE BARREN** and other adjuvants

## APPLICATION RATES

Application rate of tank mix by air or ground (gallons per acre)	Application rate of ELITE BARREN per acre
0-5	6 to 16 ounces*
5-30	1 to 4 pints
Above 30	1% minimum in spray mix

\*Ensure 1% minimum is added to the spray tank but do not exceed 5%.

## STORAGE AND DISPOSAL

**Storage:** Store in a cool well-ventilated place. Keep in original container tightly closed. Do not reuse empty container. Do not store with food, feed, or other material to be used or consumed by humans or animals. Do not contaminate water supplies. For optimal storage, store between 40° and 90° F.

**Disposal:** Do not contaminate water, food or feed by disposal. Dispose of contents in container to an approved waste disposal facility in accordance with all local, regional, national and international regulations. Triple rinse (or equivalent) adding rinse water to application tank. Offer container for recycling or dispose of in a sanitary landfill or by other procedures approved by local regulations. The Agricultural Container Recycling Council (ACRC) operates the national recycling program. To contact your state and local ACRC recycler visit the ACRC web page at [www.acrcycle.org](http://www.acrcycle.org).

## WARRANTY

Red River warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to the label instruction, or any abnormal conditions, or under conditions not reasonably foreseeable to Red River, and buyer assumes the risk of any such use.

Manufactured for:

