Seal No Copy

Rate of 3336 DG Lite Remarks *

For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

Table 3: Turf Disease Control

Colletotrichum graminicola

Bermudagrass Decline

Cool Season Brown Pato

Gaeumannomyces graminis vai

Anthracnose, basal

Take-All-Patch

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMA
CAUTION

CAUTION: Harmful if swallowed, absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. PERSONAL PROTECTIVE EQUIPMENT (PPE) for Category C on the EPA chemical resistance selection char

 Chemical resistant gloves are required for applicators using hand held equipment. USER SAFETY REQUIREMENTS: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately i pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove clothing/PPE immediately after handling this product As soon as possible, wash thoroughly and change into clean clothing. ENVIRONMENTAL HAZARDS

Do not apply directly to water, or areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water. DIRECTIONS FOR USE
IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. For use only by certified applicators

and those under their immediate supervision. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any other requirements specific to your State or Tribe, consult the agency

AGRICULTURAL USE REQUIREMENTS Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It

Long-sleeved shirt and long pants

box apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours Personal Protective Equipment (PPE) required for early entry to treated areas that is permitted under the Worker Protection Standard and that

· Coveralls over long-sleeved shirt and long pants Chemical-resistant gloves made of any waterproof materia Chemical-resistant footwear plus socks

NON-AGRICULTURAL USE REQUIREMENTS The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural Turf and Landscape Uses: Do not enter or allow others to enter until dusts have settled.

STORAGE AND DISPOSAL

Pesticide Storage: Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, contain/re-capture spillage and use in accordance with Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. I these wastes cannot be used according to the label instructions, contact your state Pesticide or Environmental Control Agency, or the Hazardous

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning.

COMMERCIAL TURF AND ORNAMENTA

Chemical-resistant headgear for over head exposures

3336 DG Lite is a broad spectrum fungicide exhibiting preventative, curative and systemic properties. It is useful on a wide variety of turf and ornamental disease problems. Apply 3336 DG Lite with ground based, hand operated (push) or mechanical (power) equipment sufficient to accurately deliver this product to the application site. Do not apply by hand or with hand held rotary spreader (e.g. bellygrinder). Application equipment should be properly calibrated prior to use. Application equipment should be thoroughly cleaned after use. If a rate range is given, use the higher rate under conditions of severe disease pressure. Contact your local State Extension service for recommendations for application schedules appropriate to your area. This product is not for homeowner use and can only be used by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products. No claim of compatibility with other pesticides is implied. Do not apply this product through an

Resistance Management: To avoid the development of tolerant strains of fungi, 3336 DG Lite should be used with fungicides of different mode of action. Cleary Chemical does not recommend the use of products containing thiabendazole in combination or rotation with 3336 DG Lite. These the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultura fungicide or disease control strategy. As long as recommended precautions are followed, 3336 DG Lite can remain useful for disease control. COMMERCIAL TURF APPLICATIONS

centers), and golf courses (greens, tees, fairways and aprons) of cool and warm season grasses such as Bentgrass, Bluegrass, Bermudagrass Fescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures. 3336 DG Lite is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. 3336 DG Lite is to be used for the prevention and control of the diseases mentioned below. 3336 DG Lite ma be applied to dry or moist leaf/soil conditions and should be activated with irrigation or natural rainfall within 24 hours for best results. If irrigated application should be watered in by applicator with 0.1-0.25 inches of water to begin activation of product for disease control. When treatin golf greens, treatment of aprons is recommended. Use the highest recommended rate under conditions of severe disease pressure. Do not graze nimals on treated turf. Do not feed clippings to livestock or poultry. Not for use on turf being grown for sale or other commercial use as sod. Table 1: Maximum Individual Application Rates and Minimum Retreatment Intervals Do not exceed the amounts per acre or reduce the re-treatment interval indicated below

Use Site	Maximum application rate of 3336 DG Lite	Minimum Retreatment Interval	Comments	
Residential or Public Areas	131 lb/Acre (3 lb / 1,000 sq ft)	14 days		
Golf Course Tees, Greens, Aprons	392 lb/Acre (9 lb / 1,000 sq ft)	14 days		
Golf Course Fairways –except Florida	262 lb/Acre (6 lb / 1,000 sq ft)	14 days	Excludes Florida	
Golf Course Fairways – Florida Only	131 lb/Acre (3 lb / 1,000 sq ft)	14 days	Florida Only	
Table 2: Maximum Seasonal Appl Do not exceed the following amounts				
Use Site	Maximum Pounds 3336 DG Lite per Acre per Season	Pounds 3336 DG Lite per 1,000 sq ft	Comments	
Residential or Public Areas	523 lb	12 lb	4 Applications per ye	
Golf Course Tees, Greens, Aprons	1048 lb	24 lb		
Golf Course Fairways –except Florida	262 lb	6 lb	Excludes Florida	
Golf Course Fairways – Florida Only	131 lb	3 lb	Florida Only	

For prevention in historic areas of disease pressure, apply twice at 14 day intervals when soil temperature reaches 60°F. For curative control, apply Apply in mid-July or when disease symptoms first appear and repeat at 14 day intervals for suppression. Use higher rates under most severe disease expression. Follow proper agronomic recommendations to maintain plant vigor For prevention, apply in fall before turf has stopped all growth activity. Apply second application in early Spring when soil temperatures reach 55-60°I appears in early spring and continue at 14 day intervals. Apply 2 treatments at 21 day intervals in late fall to early winter, with the last application made just prior to first permanent snow cover. Rotations with PCNB For prevention, use lowest rate and apply application before expected period of disease development. Apply when disease first appears and continue at 14 day intervals. Rotations with chlorothalonil, iprodione, or mancozeb (Protect™)

Fusarium Blight Apply when disease first appears at 14 day intervals Gray Leaf Spot (Blast) ntinue applications at 14 day intervals as needed. s needed. Rotations with chlorothalonil, iprodione, or mancozeb (Protect) are Leaf, crown, and root disease Bipolaris, Curvularia, and Pink Snow Mold Apply middle-late Fall to early Winter before turf has stopped all growth activity Michrodochium nivale or in spring during periods of thaw, or when disease symptoms first appear For suppression, apply at 14 day intervals when disease first appears. ecommendations. Apply at 14 day intervals as needed. Rotations and/or tank

can be utilized.

*Observe the maximum individual application rates and maximum seasonal application limits in Table 2.							
Table 4: Guide to 3336 DG Lite Re These settings are presented as a Spreader calibration at the beginni	guide for use. Age	e, condition of s	preader and or		se variation in applied	d rates of materia	
		Swath (FT)		Pounds of 3336 DG Lite Applied per 1,000 Square Feet			
Spreader	Speed (MPH)		Cone Set	3	6	9	
Lesco Rotary	3	8	0.25	13	17	21	
Cyclone Rotary	3	4	n/a	4	5 1/4	6 ½	
Scotts R-8A1 Rotary	3	11	6	J 1/2	N	Q	
The Andrease Assubra 0000	0		_	1/ 1/	0		

Lesco Rotary	3	8	0.25	13	17	21
Cyclone Rotary	3	4	n/a	4	5 1/4	6 ½
Scotts R-8A1 Rotary	3	11	6	J ½	N	Q
The Andersons AccuPro 2000	3	11	6	K ½	0	R
The Andersons SR 2000	3	11	6	K ½	0	R
Spyker	3	5	n/a	3 5/7	4 1/2	5 1/5
PrizeLAWN CRB III & Big Foot	3	5	n/a	l 1/2	M	Р
Earthway	3	7	n/a	12	15.5	18.5
Precision Broadcast	3	4	n/a	7 3/4	10 1/4	13 ½
Precision Drop	3	n/a	n/a	5 3/4	8 1/4	11 ½
Lely W (2)*	4.5	34	III-C	5	6 3/4	8 1/4
Vicon 03 Series*	4.5	30	0	20	31	40

Nursery, Greenhouse, Landscape & Interiorscape
Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs 3336 DG Lite is a broad spectrum systemic fungicide which controls a variety of foliar, stem, and root diseases on a wide range of commercially important plants. 3336 DG Lite may be used for disease control and prevention programs on annual and perennial flowers, bedding plants, foliage plants, ground covers, and deciduous and evergreen trees and shrubs found in Nursery, Greenhouse, Interiorscape, and Landscape sites. Follow

3336 DG Lite may be used as a broadcast, fly-on, surface, over the top, and media incorporation application for the diseases listed below. For surface and broadcast applications to established plants apply at the rates below. For cut flowers do not exceed the maximum application rate 0.5 o ai/acre/application. 3336 DG Lite may be applied to dry or moist soil conditions and should be activated with irrigation or natural rainfall within 4 hours for best results. If irrigated, application should be watered in by applicator with 0.1-0.25 inches of water to begin activation. For media ncorporation, mix at a rate of 2-3 lb/cu yd before seeding or transplanting of desired crop. Media incorporated with 3336 DG Lite may be stockpile Note: The "Directions For Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any

For control of soil borne diseases effecting

All other trademarks are the property of their respective owners.

.5' Seal No Copy

Bulbs, Corms, Rhizomes, Roots, Crowns Botrytis, Cylindrocladium, Fusarium, Gliocladium Myrothecium, Penicillium, Ramularia,

transplanting to propagation beds, containers, pots, or nursery or landscape beds. For curative control, apply after seeding or transplanting when disease first appears. Repeat every 21-28 days for adequate crop and dipotassium salts of phosphorous acid or fosetyl-Al are required for

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOOD DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTOR' OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAI PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE FOR SPECIAL, INCIDENTAL, OR ONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION HANDLING. AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER DEMANDS WHATSOEVER. RESULTING FROM OR BY REASON OF OR RISING OUT OF THE MISUSE. OR FAILURE TO FOLLOW LABEL WARNINGS OR USER. OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. BUYER'S OR USER'S EXCLUSIVE REMEDY. AND MANUFACTURER'S If you do not agree with or do not accept any of the directions for use, the warranty disclaimers, or limitations of liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Nufarm Grow a better tomorrow.

plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This

3356 DGLITE®

Turf and Ornamental Systemic Fungicide

For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

ACTIVE INGREDIENT

	TOTAL	100.00%
OTHER INGREDIENTS		97.92%
Thiophanate-methyl		2.08%

Keep Out Of Reach Of Children CAUTION

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

FIRST AID

IF SWALLOWED

Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. You may also contact (877) 325-1840 for emergency medical treatment information.

EPA Reg. No. 1001-70 EPA Est. No. 9198-0H-1M, 9198-0H-2B, 9198-AL-001A **Underlined letter is first** letter used in run code on bag

Net Contents

| Manufactured for: Cleary Chemicals LLC 11901 S. Austin Avenue Alsip, IL 60803



10495032



.5' Seal No Copy

.5' Seal No Copy

.5' Seal No Copy

.5' Seal No Copy