



FINESSE®

CEREAL AND FALLOW HERBICIDE

*For Use on Wheat, Barley, Triticale and Fallow
Dry Flowable*

CHLORSULFURON	GROUP	2	HERBICIDE
METSULFURON-METHYL	GROUP	2	HERBICIDE

<i>Active Ingredient</i>	<i>By Weight</i>
Chlorsulfuron	62.5%
Metsulfuron Methyl	12.5%
Other Ingredients	25.0%
TOTAL	100%

EPA Reg. No. 279-9610

EPA Est. No. 352-IL-001

U.S. Pats. 4,127,405 & 4,383,113

Contains 0.75 lb. active ingredient per pound (chlorsulfuron 0.625 and metsulfuron methyl 0.125 lb. active ingredient per pound).

Nonrefillable Container

Refillable Container

Net: Weight 20 oz

OR

Net: _____

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center, doctor or going for treatment.

You may also contact 1-800-331-3148 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco or using the toilet. Avoid breathing dust or spray mist. Remove and wash contaminated clothing before reuse.

For medical emergencies involving this product, call toll free 1-800-331-3148.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders, applicators, and other handlers must wear:

Long-sleeved shirt and long pants.

Chemical Resistant Gloves made of any waterproof material such as polyethylene or polyvinyl chloride.

Shoes plus socks

Discard clothing and other absorbent material that have been drenched or heavily contaminated with this product. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

Sold By



FMC Corporation
2929 Walnut Street
Philadelphia, PA 19104

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

Groundwater Advisory

This product is known to leach through soil into groundwater under certain conditions as a result of label use. This product may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of this product from runoff water and sediment. Runoff of this product will be greatly reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

Windblown Soil Particles Advisory

This product has the potential to move off-site due to wind erosion. Soils that are subject to wind erosion usually have a high silt and/or fine to very fine sand fractions and low organic matter content. Other factors which can affect the movement of windblown soil include the intensity and direction of prevailing winds, vegetative cover, site slope, rainfall, and drainage patterns. Avoid applying this product if prevailing local conditions may be expected to result in off-site movement.

Non-target Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated area. Protect the forage and habitat of non-target organisms by minimizing spray drift. For further guidance and instructions on how to minimize spray drift, refer to the Spray Drift Management section of this label.

PESTICIDE HANDLING

- Calibrate sprayers only with clean water away from the well site.
- Make scheduled checks of spray equipment.
- Ensure that all operation employees accurately measure pesticides.
- Mix only enough product for the job at hand.
- Avoid over-filling of spray tank.
- Dilute and agitate excess solution and apply at labeled rates or uses.
- Avoid storage of pesticides near well sites.
- When triple rinsing the pesticide container, be sure to add the rinsate to the spray mix.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

FINESSE® cereal and fallow herbicide, referred to below as FINESSE herbicide, must be used only in accordance with instructions on this label, or as otherwise permitted by FIFRA.

Always read the entire label, including the Limitation of Warranty and Liability.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls.

Chemical Resistant Gloves made of any waterproof material.

Shoes plus socks.

To the extent consistent with applicable law, FMC will not be responsible for losses or damages resulting from the use of this product in any manner not specifically directed by FMC.

PRODUCT INFORMATION

FINESSE herbicide is registered for use on land primarily dedicated to the long-term production of wheat, barley, or triticale.

FINESSE herbicide is a dry-flowable granule that controls weeds in wheat (including durum), barley, triticale and fallow.

FINESSE herbicide is mixed in water or may be slurried in water then added directly into liquid nitrogen fertilizer solutions and applied as a uniform broadcast spray. A surfactant needs to be used in the spray mix unless otherwise specified on this label.

FINESSE herbicide is noncorrosive, nonflammable, nonvolatile, and does not freeze.

FINESSE herbicide controls weeds by both preemergence and postemergence activity. For best preemergence results, apply FINESSE herbicide before weed seeds germinate. Use sprinkler irrigation or allow rainfall to move FINESSE herbicide 2" to 3" deep into the soil profile.

For best postemergence results, apply FINESSE herbicide to young, actively growing weeds. The use rate depends upon the weed spectrum and size of weeds at the time of application. The degree and duration of control may depend on the following:

- weed spectrum and infestation density
- weed size at application
- environmental conditions at and following treatment

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

FINESSE herbicide is absorbed through the roots and foliage of plants, rapidly inhibiting the growth of susceptible weeds. For preplant and preemergence weed control, rainfall is needed to move FINESSE herbicide into the soil. Weeds will generally not emerge from preplant and preemergence applications. In some cases, susceptible weeds may germinate and emerge a few days after application, but growth then ceases, and leaves become chlorotic three to five days after emergence. Death of leaf tissue and growing point will follow in some species, while others will remain green but stunted and noncompetitive.

One to three weeks after postemergence application to weeds, leaves of susceptible plants appear chlorotic, and the growing point subsequently dies. In warm, moist conditions, the expression of herbicide symptoms is accelerated; in cold, dry conditions, expression of herbicide symptoms is delayed. Death of leaf tissue will follow in some species, while others will remain green but stunted and noncompetitive. Postemergence weed control may be reduced if rainfall occurs within 6 hours after application.

FINESSE herbicide provides the best control of weeds in vigorously growing crops that shade competitive weeds. Weed control in areas of thin crop stand or seeding skips may not provide satisfactory control. However, a crop canopy that is too dense at application can intercept spray and reduce weed control.

The herbicidal action of FINESSE herbicide may be less effective on weeds stressed from adverse environmental conditions (such as extreme temperatures or moisture, drought stress), abnormal soil conditions, or cultural practices that increase weed stress. In these cases, tank mix FINESSE herbicide with other registered herbicides (including 2,4-D or MCPA) to aid in control. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflicts with this FINESSE herbicide label, DO NOT use in a tank mixture with FINESSE herbicide.

RESTRICTIONS

- Do not apply to frozen ground where surface runoff may result.
- Do not apply to snow-covered ground.
- Do not discharge excess material on the soil at a single spot in the field or mixing/loading station.
- Do not apply to irrigated land where tailwater will be used to irrigate other cropland.
- Do not use in Alamosa, Conejos, Costilla, Rio Grande, and Saguache counties of Colorado.

• Do not use less than 0.2 oz per acre of FINESSE herbicide preplant, preemergence or postemergence	FINESSE herbicide	Active Ingredient Equivalent	
	Rate Oz/A	Chlorsulfuron Lb. ai/A	Metsulfuron-methyl Lb. ai/A
	0.2	0.0078	0.0016

- Wherever FINESSE herbicide is used on land previously treated with Glean® XP herbicide, Ally® XP herbicide, Ally® Extra SG herbicide (with TotalSol® soluble granules), Amber® Custom-Pak™ herbicide or other longer residual herbicides with the same mode of action, you must read the rotational guidelines on both labels and follow the one with the longest interval stated for your situation before choosing to rotate to crops other than wheat or barley.
- To reduce the potential for movement of treated soil due to wind erosion, do not apply to powdery, dry, or light sandy soils until they have been stabilized by rainfall, trashy mulch, reduced tillage or other cultural practices. Injury to adjacent crops may result when treated soil is blown onto land used to produce crops other than cereal grains.
- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
 - Do not apply, drain, or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.
 - Do not use on lawns, walks, driveways, or tennis courts.
- For all applications with products containing these active ingredients do not exceed 0.0625 oz ai/A (Lb. ai/A 0.0039) of metsulfuron methyl and 0.37 oz ai/A (Lb. ai/A 0.0231) of chlorsulfuron in a year.

PRECAUTIONS

- Injury to or loss of desirable trees or vegetation may result from failure to observe the following:
 - Take all necessary precautions to avoid all direct or indirect contact (such as spray drift) with non-target plants or areas.
 - Carefully observe sprayer cleanup instructions, both prior to and after using this product, as spray tank residue may damage crops other than wheat or barley.

WEED RESISTANCE MANAGEMENT

FINESSE herbicide, which contains the active ingredients chlorsulfuron and metsulfuron methyl is a group 2 herbicide based on the mode of action classification system of the Weed Science Society of America.

Proactively implementing diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides is a best practice. A diversified weed management program may include the use of multiple herbicides with different sites of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and directions for use is important to delay the selection for resistance.

The continued effectiveness of this product depends on the successful implementation of a weed resistance management program.

To aid in the prevention of developing weeds resistant to this product, users should:

- Scout fields before application to ensure herbicides and rates will be appropriate for the weed species and weed sizes present.
- Start with a clean field, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small (less than 4 inches).
- Apply full rates of FINESSE herbicide, for the most difficult to control weed in the field at the specified time (correct weed size) to minimize weed escapes.
- Scout fields after application to detect weed escapes or shifts in control of weed species.
- Control weed escapes before they reproduce by seed or proliferate vegetatively.
- Report any incidence of non-performance of this product against a particular weed to your FMC representative, local retailer, or county extension agent.
- Contact your FMC representative, crop advisor, or extension agent to find out if suspected resistant weeds to these MOAs have been found in your region. Do not assume that each listed weed is being controlled by multiple sites of action. Products with multiple active ingredients are intended to broaden the spectrum of weeds that are controlled. Some weeds may be controlled by only one of the active ingredients in this product.
- If resistance is suspected, treat weed escapes with an herbicide having a site of action other than Group 2 and/or use nonchemical methods to remove escapes, as practical, with the goal of preventing further seed production.

- Suspected herbicide-resistant weeds may be identified by these indicators:
 - Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
 - A spreading patch of non-controlled plants of a particular weed species; and
 - Surviving plants mixed with controlled individuals of the same species

Additionally, users should follow as many of the following herbicide resistance management practices as is practical:

- Use a broad spectrum soil-applied herbicide with other sites of action as a foundation in a weed control program.
- Utilize sequential applications of herbicides with alternative sites of action.
- Rotate the use of this product with non-Group 2 herbicides.
- Avoid making more than two applications of FINESSE herbicide, and any other Group 2 herbicides within a single growing season unless mixed with an herbicide with a different site of action with an overlapping spectrum for the difficult-to- control weeds.
- Incorporate non-chemical weed control practices, such as mechanical cultivation, crop rotation, cover crops and weed-free crop seeds, as part of an integrated weed control program.
- Use good agronomic principles that enhance crop development and crop competitiveness.
- Thoroughly clean plant residues from equipment before leaving fields suspected to contain resistant weeds.
- Manage weeds in and around fields, during and after harvest to reduce weed seed production.

INTEGRATED PEST MANAGEMENT

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

CEREALS APPLICATIONS

Precautions:

- Wheat, barley, and triticale varieties may differ in their response to various herbicides. FMC advises that you first consult your state experiment station, university, or extension agent as to sensitivity to any herbicide. If no information is available, limit the initial use of FINESSE herbicide to a small area.
- Temporary discoloration and/or crop injury may occur if FINESSE herbicide is applied when the crop is stressed by severe weather conditions (such as heavy rainfall, prolonged cold weather, or wide fluctuations in day/night temperatures), disease or insect damage, low fertility, applications to coarse soils, or when applied in combination with surfactant and high rates of liquid nitrogen fertilizer solutions.

PREPLANT AND PREEMERGENCE

FINESSE herbicide can be tank mixed with other products registered for preplant/preemergence use in wheat including products with the active ingredient glyphosate. In the case of tank mixes with other herbicides, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this FINESSE herbicide label, DO NOT use in a tank mixture with FINESSE herbicide.

Restrictions:

- Do not apply FINESSE herbicide preplant or preemergence on durum or Wampum wheat, barley, or triticale.
- Do not apply preemergence or preplant incorporated to late fall plantings when cold and/or dry weather can cause delayed seedling emergence and/or stress to seedling plants. Under these conditions, wait until crop has emerged and is showing good vigor before making a postemergence treatment.

Precautions:

- Crop injury may result when preemergence or preplant incorporated applications of FINESSE herbicide are made to wheat seeded less than 1" deep.
- Crop injury may result if FINESSE herbicide is used where an organophosphate insecticide has been applied or is intended for use as an in-furrow treatment.

APPLICATION INFORMATION	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
<p>Winter Wheat</p> <p><u>Preplant:</u> Apply FINESSE herbicide at 0.2 to 0.5 oz per acre (before winter wheat is planted). In TX, OK, KS, NE, and SD, preplant application at 0.2 to 0.5 may be shallow incorporated into the top 1 inch of soil.</p> <p><u>Preemergence:</u> Apply FINESSE herbicide at 0.2 to 0.5 oz per acre (after planting but before winter wheat emerges). In WY, MT, ND and MN, DO NOT exceed 0.3 oz per acre preemergence.</p> <p>Spring Wheat</p> <p><u>Preplant/Preemergence:</u> Apply FINESSE herbicide at 0.2 to 0.4 oz per acre in spring wheat (except Durum wheat and Wampum variety of Spring Wheat). In WY, MT, ND, SD, and MN, DO NOT exceed 0.3 oz per acre preplant or preemergence.</p>	0.2	0.0078	0.0016
	0.3	0.0117	0.0023
	0.4	0.0156	0.0031
	0.5	0.0195	0.0039

Restrictions:

Crop/Use	Application Timing	Active Ingredient	Maximum Oz/A of FINESSE herbicide per Single Application	Maximum Lb. ai/A per Single Application	Maximum Oz/A of FINESSE per Year	Maximum Lb.ai/A per Year	Maximum Number of Applications per Year	Minimum Treatment Interval (Days)	Pre-Harvest Interval, Days
Winter Wheat	Preplant (before winter wheat is planted)	Chlorsulfuron	0.5	0.0195	0.5	0.0195	1	Not Applicable	45(for grain)
		Metsulfuron-methyl		0.0039		0.0039			
Winter Wheat (TX, OK, KS, NE, & SD)	Preplant (may be shallow incorporated into the top 1 inch of soil)	Chlorsulfuron	0.5	0.0195	0.5	0.0195	1	Not Applicable	45(for grain)
		Metsulfuron-methyl		0.0039		0.0039			
Winter Wheat	Preemergence (After planting but before winter wheat emerges)	Chlorsulfuron	0.5	0.0195	0.5	0.0195	1	Not Applicable	45(for grain)
		Metsulfuron-methyl		0.0039		0.0039			
Winter Wheat (WY, MT, ND, & MN)	Preemergence	Chlorsulfuron	0.3	0.0117	0.3	0.0117	1	Not Applicable	45(for grain)
		Metsulfuron-methyl		0.0023		0.0023			
Spring Wheat (Except Durum and Wampum varieties)	Preplant or Preemergence	Chlorsulfuron	0.4	0.0156	0.4	0.0156	1	Not Applicable	45(for grain)
		Metsulfuron-methyl		0.0031		0.0031			
Spring Wheat (WY, MT, ND, SD, & MN)	Preplant or Preemergence	Chlorsulfuron	0.3	0.0117	0.3	0.0117	1	Not Applicable	45(for grain)
		Metsulfuron-methyl		0.0023		0.0023			

- Do not apply FINESSE herbicide preemergence on cereals if the seed has germinated and has started to emerge above the soil surface.
- Do not use FINESSE herbicide preemergence on cereals that have been planted into dry soil (“dusted in”) or on very coarse, uneven seedbeds

POSTEMERGENCE

FINESSE herbicide may not be used within 60 days of crop emergence if an organophosphate insecticide was used as an in-furrow treatment or crop injury may result.

In areas where late fall or winter cold weather conditions are unpredictable and can be severe (such as the Pacific Northwest and Northern plains), to avoid crop injury due to cold weather, DO NOT make applications during the 1 to 4-leaf stage of wheat, barley, or triticale. The combined effects of herbicide stress plus cold weather stress can result in greater crop injury than either stress factor alone.

APPLICATION INFORMATION	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
<u>Postemergence:</u> Apply FINESSE herbicide at 0.2 to 0.4 oz per acre to wheat or barley any time after the crop is in the 1-leaf stage, but before boot stage. Apply FINESSE herbicide at 0.2 to 0.4 oz per acre to triticale any time after the crop is in the 2-3 leaf stage but before the flag leaf is visible.	0.2	0.0078	0.0016
	to 0.4	to 0.0156	to 0.0031

DO NOT apply FINESSE herbicide during the boot stage or early heading stage, as crop injury may result.

Restrictions:

Crop/Use	Application Timing	Active Ingredient	Maximum Oz/A of FINESSE herbicide per Single Application	Maximum Lb. ai/A per Single Application	Maximum Oz/A of FINESSE per Year	Maximum Lb.ai/A per Year	Maximum Number of Applications per Year	Minimum Treatment Interval (Days)	Pre-Harvest Interval, Days
Wheat & Barley	Postemergence (After 1-leaf stage but before boot stage)	Chlorsulfuron	0.4	0.0156	0.4	0.0156	2	14	45(for grain)
		Metsulfuron-methyl		0.0031		0.0031			
Wheat, Barley, Triticale	Postemergence (Pacific Northwest & Northern Plains) After the 4-leaf stage but before boot stage	Chlorsulfuron	0.4	0.0156	0.4	0.0156	1	Not Applicable	45(for grain)
		Metsulfuron-methyl		0.0031		0.0031			
Triticale (Areas other than Pacific Northwest & Northern Plains)	Postemergence (any time after the crop is in the 2-3 leaf stage but before the flag leaf is visible)	Chlorsulfuron	0.4	0.0156	0.4	0.0156	1	Not Applicable	45(for grain)
		Metsulfuron-methyl		0.0031		0.0031			

- Do not apply to wheat, barley, or triticale undersown with legumes and grasses, as injury to the forages will result.
- Wherever FINESSE herbicide is used on land previously treated with Glean® XP herbicide, Ally® XP herbicide, Ally® ExtraSG herbicide (with Totalso® soluble granules), Amber® Custom-Pak™ herbicide or other longer residual herbicides with the same mode of action, read the rotational guidelines on both labels and follow the one with the longest interval stated for your situation before choosing to rotate to crops other than wheat or barley

Precautions:

- For ground applications applied postemergence to weeds when dry, dusty field conditions exist, control of weeds in wheel track areas may be reduced. The addition of 2,4-D or MCPA could improve weed control under these conditions.

FALLOW APPLICATIONS

APPLICATION INFORMATION	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
FINESSE herbicide may be used as a fallow treatment and may be tank mixed with other herbicides that are registered for use in fallow. Apply FINESSE herbicide at 0.2 – 0.4 oz per acre in the spring through the fall when the majority of weeds have emerged and are actively growing.	0.2	0.0078	0.0016
	to 0.4	to 0.0156	to 0.0031

In the case of tank mixes with other herbicides, it is the pesticide user’s responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this FINESSE herbicide label, DO NOT use in a tank mixture with FINESSE herbicide.

Restrictions:

Crop/Use	Application Timing	Active Ingredient	Maximum Oz/A of FINESSE herbicide per Single Application	Maximum Lb. ai/A per Single Application	Maximum Oz/A of FINESSE per Year	Maximum Lb.ai/A per Year	Maximum Number of Applications per Year	Minimum Treatment Interval (Days)	Pre-Harvest Interval, Days
Fallow	-----	Chlorsulfuron	0.4	0.0156	0.4	0.0156	2	1	-----
		Metsulfuron-methyl		0.0031		0.0031			

BORDER AREA APPLICATIONS

APPLICATION INFORMATION	FINESSE® herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
FINESSE herbicide may be used for control of broadleaf weeds in field border areas and fence lines. Apply FINESSE herbicide at 0.2 to 0.5 oz per acre.	0.2	0.0078	0.0016
	to	to	to
	0.5	0.0195	0.0039

Restrictions:

Crop/Use	Application Timing	Active Ingredient	Maximum Oz/A of FINESSE herbicide per Single Application	Maximum Lb. ai/A per Single Application	Maximum Oz/A of FINESSE per Year	Maximum Lb.ai/A per Year	Maximum Number of Applications per Year	Minimum Treatment Interval (Days)	Pre-Harvest Interval, Days
Border Area (Field border areas & Fence Lines)	-----	Chlorsulfuron	0.5	0.0195	0.5	0.0195	2	1	-----
		Metsulfuron-methyl		0.0039		0.0039			

SURFACTANTS - ALL CROPS

Unless otherwise specified, add a nonionic surfactant having at least 80% active ingredient at 0.125 to 0.5% v/v (0.5 to 2 qt per 100 gal of spray solution).

The higher rate of surfactant is particularly effective with spray volumes of 5 gallons per acre (GPA) or less and when using low rates of FINESSE herbicide. Consult your agricultural dealer, applicator, or FMC representative for a listing of specified surfactants.

Antifoaming agents may be used if needed.

DO NOT use low rates of liquid nitrogen fertilizer solution as a substitute for surfactant.

WEEDS CONTROLLED

FINESSE herbicide effectively controls the following weeds when applied at the rates shown:

APPLICATION INFORMATION	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
0.2 to 0.3 oz per acre	0.2	0.0078	0.0016
	to	to	to
	0.3	0.0117	0.0023

Blue mustard	Mayweed chamomile
Broadleaf dock	Miners lettuce
Bur beakchervil	Pineappleweed
Bur buttercup (testiculate)	Prickly lettuce††
Carolina geranium	Prostrate pigweed
Chickweed (common‡, jagged, mouseear)	Plains coreopsis
Conical catchfly	Purslane
Corn spurry	Redstem filaree
Cow cockle	Redroot pigweed‡
Curly dock	Sheperd's purse
Cutleaf evening primrose	Smallseed falseflax‡
False chamomile	Smooth pigweed‡
Field pennycress	Tansymustard*†
Flixweed*†‡	Treacle mustard
Groundsel	(Bushy wallflower)
Hempnettle	Tumble mustard (Jim Hill)
Henbit	Virginia pepperweed
Lady's thumb	White cockle
Lambsquarters‡	Wild mustard‡
	Wild carrot

APPLICATION INFORMATION	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
0.3 to 0.4 oz per acre	0.3	0.0117	0.0023
	to	to	to
	0.4	0.0156	0.0031

Annual bluegrass*†‡	Kochia*†‡
Annual ryegrass*†‡	Pennsylvania smartweed*
Annual sowthistle	Persian darnel*†
Bedstraw*†	Prickly poppy (pinnate)
Bromus species (cheat, Downy brome, Japanese brome)*†	Russian thistle*†‡
Canada thistle*†	Speedwell (common, ivyleaf)*
Coast fiddleneck (tarweed)	Sunflower†‡
Corn groomwell*†	Vetch†
Dove foot geranium	Volunteer corn†
Green foxtail (pigeongrass)*‡	Wild buckwheat†
Knotweed (prostrate)*†	Wild radish†
	Yellow foxtail*†‡

APPLICATION INFORMATION	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
0.5 ounces per acre (prior to winter wheat emergence only)	0.5	0.0195	0.0039

Annual ryegrass*†‡
Bromus species (cheat, downy brome, Japanese brome)*†‡
Volunteer corn†

* When used as directed, weeds are suppressed and/or controlled. Weed suppression is a visible reduction in weed competition (reduced population and/or vigor) as compared to an untreated area. Degree of suppression will vary with rate used, size of weeds, and environmental conditions following treatment.

† See the **Specific Weed Instructions** section for more information regarding controlling and suppressing these weeds.

‡ Naturally occurring resistant biotypes of these weeds are known to occur. See **Tank Mixtures, Specific Weed Instructions, and Weed Resistance** sections of this label for additional information.

SPECIFIC WEED INSTRUCTIONS

Annual bluegrass/annual ryegrass

FINESSE herbicide Preplant or Preemergence

Apply FINESSE herbicide at 0.5 oz per acre preplant or after planting winter wheat but before wheat emerges. or Apply FINESSE herbicide at 0.5 oz per acre preplant or after planting winter wheat but before wheat emerges followed by a sequential application of metribuzin in the fall once the wheat has reached the 4 to 5-leaf stage of growth and the annual grassy weeds are in the 1 to 3-leaf stage of growth. Refer to the product containing the active ingredient metribuzin label for specific rates and use instructions. or For improved control in the Pacific Northwest, apply a tank mix of FINESSE herbicide at 0.3 to 0.4 oz per acre plus Karmex® DF herbicide at labeled rates preemergence to bluegrass or ryegrass. One-half to 1" of rainfall is needed to move the herbicides into the weed root zone prior to bluegrass or ryegrass emergence. Refer to the product containing the active ingredient diuron label for specific rates and use instructions.	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
	0.3	0.0117	0.0023
	0.4	0.0156	0.0031
	0.5	0.0195	0.0039

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this FINESSE herbicide label, DO NOT use in a tank mixture with FINESSE herbicide.

FINESSE herbicide Postemergence

In the case of tank mixes with other herbicides, it is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each

product in the tank mixture. If the instructions on the tank mix partner label conflict with this FINESSE herbicide label, DO NOT use in a tank mixture with FINESSE herbicide.

<p>Apply a tank mix of FINESSE herbicide at 0.2 to 0.4 oz per acre and metribuzin at labeled rates postemergence to the crop and grassy weeds when wheat has reached the 4 to 5-leaf stage of growth and the grassy weeds have reached the 1 to 3-leaf stage of growth. Refer to the product containing the active ingredient metribuzin for rates and specific use instructions.</p> <p>Note: See Bromus species (cheat, downy brome, Japanese brome) section for additional information on the use of metribuzin.</p> <p>Bedstraw: Apply FINESSE herbicide at 0.4 oz per acre. For postemergence treatments, apply before bedstraw is over 2" long; use 2 qt of surfactant per 100 gal of spray solution.</p>	FINESSE herbicide		Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)	
	0.2	0.0078	0.0016	
0.4	0.0156	0.0031		

Bromus species (cheat, downy brome, Japanese brome): Best suppression of these grasses is achieved by applications of FINESSE herbicide with metribuzin either in tank mixtures or as sequential treatments.

Additional information may be available in a metribuzin supplemental label for winter wheat, barley, and fallow.

Allow for adequate rainfall (1/2 to 1") to move FINESSE herbicide and metribuzin into the weed root zone before weeds germinate and develop an established root system. Lack of adequate rainfall following application will result in reduced performance.

To avoid the risk of cold weather-related crop injury and lack of performance, apply metribuzin before winter dormancy of the crop and grassy weeds. Excessive rainfall immediately after application may result in crop injury. DO NOT tank mix FINESSE herbicide plus metribuzin with any other pesticide other than surfactants specified on either the FINESSE herbicide or metribuzin labels. Apply only to metribuzin-approved varieties, see metribuzin label for listing of sensitive wheat and barley varieties.

Preemergence/Sequential Applications

<p>Apply FINESSE herbicide at 0.5 oz per acre preemergence after planting winter wheat but before wheat emerges. A sequential application of metribuzin may be applied at label rates in the fall once the wheat has reached the 4 to 5-leaf stage of growth and the annual grassy weeds are in the 1 to 3-leaf stage of growth. Refer to the product containing the active ingredient metribuzin for rates and specific use instructions.</p> <p><u>Idaho, Oregon, and Washington</u>—Apply FINESSE herbicide at 0.4 to 0.5 oz per acre after planting winter wheat but before wheat emerges.</p>	FINESSE herbicide		Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)	
	0.4	0.0156	0.0031	
0.5	0.0195	0.0039		

If suppression of brome grass is not satisfactory following the preemergence application of FINESSE herbicide, apply a sequential treatment of metribuzin at lower rates in the fall when the crop is in the 2-leaf to 3 tiller stage or at higher rates after winter wheat has at least 4 tillers, 2 inches of secondary root systems throughout the field and actively growing. Refer to the product containing the active ingredient metribuzin label for specific rates and use instructions.

Postemergence Tank-Mix Applications

<p>Apply a tank mix of FINESSE herbicide at 0.2 to 0.4 oz per acre and metribuzin at label rates for postemergence applications to the crop and grassy weeds when wheat has reached the 4 to 5-leaf stage of growth and the grassy weeds have reached the 1 to 3-leaf stage of growth.</p> <p><u>Idaho, Oregon, and Washington</u>—Where broadleaf weeds and brome grass are the problem, apply a tank mix of FINESSE herbicide at 0.3 to 0.4 oz per acre and metribuzin at lower rates in the fall when wheat or barley is in the 2-leaf to 3-tiller stage or use FINESSE herbicide at 0.3 to 0.4 oz and metribuzin at higher rates when wheat or barley has at least 4 tillers, 2 inches of secondary root systems throughout the field and actively growing. For best results, make application before brome grass is in the 2 to 3 leaf stage. Refer to the product containing the active ingredient metribuzin label for specific rates and use instructions.</p>	FINESSE herbicide		Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)	
	0.2	0.0078	0.0016	
0.3	0.0117	0.0023		
0.4	0.0156	0.0031		

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this FINESSE herbicide label, DO NOT use in a tank mixture with FINESSE herbicide.

Canada thistle: Apply FINESSE herbicide with surfactant after the majority of thistles have emerged and while they are small (rosette stage to 4" - 6" tall) and actively growing. For maximum long-term effect, yearly treatment may be required.

Corn groomwell: Apply FINESSE herbicide at 0.4 oz per acre and apply postemergence to the crop when weeds are small and actively growing.	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
	0.4	0.0156	0.0031

Flixweed, Tansymustard: For best results, tank mix FINESSE herbicide with 2,4-D or MCPA (esters or amines) and apply postemergence when weeds are actively growing.

Foxtail/Pigeongrass (green and yellow) (MT, ND, SD and WY): Apply FINESSE herbicide at 0.4 oz per acre in the fall or spring for suppression of these foxtail species. Application before the foxtail germinates is preferred. After emergence, best results are obtained if application is made before the foxtail is more than 1" tall or beyond the 2-leaf stage. 1/2 to 1" of rainfall is needed to move FINESSE herbicide into the weed root zone before the foxtail reaches the 3-leaf stage.	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
	0.4	0.0156	0.0031

Kochia, Russian thistle, Prickly lettuce: For best results, FINESSE herbicide may be applied postemergence in the spring. Apply when kochia, Russian thistle, and prickly lettuce are less than 2" tall or 2" across and are actively growing. Use FINESSE herbicide in a tank mix with the active ingredient dicamba (including Banvel® herbicide/Clarity® herbicide) and/or 2,4-D and 2 qt surfactant per 100 gal of spray solution.

Persian Darnel (MT, ND, SD and WY): Apply FINESSE herbicide at 0.4 oz per acre in the fall or spring for suppression of Persian darnel. Application before the Persian darnel germinates is preferred. After emergence, best results are obtained if application is made before the Persian darnel is beyond the 2-leaf stage. 1/2 to 1" of rainfall is needed to move FINESSE herbicide into the weed root zone before the Persian darnel reaches the 3-leaf stage.	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
	0.3	0.0117	0.0023
Prostrate knotweed: For best results, apply FINESSE herbicide preemergence at 0.3 to 0.4 oz per acre to knotweed in the fall. For postemergence treatments, tank mix FINESSE herbicide at 0.3 to 0.4 oz per acre with 2,4-D, MCPA, or dicamba containing products registered for this use (including Banvel herbicide/Clarity herbicide and surfactant. Apply to small, actively growing plants (no more than 4 true leaves). For maximum postemergence control, knotweed plants need to remain actively growing for 3 to 4 days following application. Refer to the product containing diuron for specific rates and use instructions.	0.4	0.0156	0.0031

Sunflower: For best results, apply FINESSE herbicide after the majority of sunflowers have emerged and are small (not more than 2" tall) and are actively growing. Add surfactant at 2 qt per 100 gal of spray solution. If FINESSE herbicide is applied preemergence, make application in early spring to allow for timely and adequate rainfall to move FINESSE herbicide into the weed root zone before weeds germinate and develop an established root system.

Note: In areas of high rainfall, fall applications may not provide adequate residual control of sunflowers.

Deep-germinating sunflowers that emerge after a spring treatment may not be controlled.

Vetch: For best results, apply FINESSE herbicide postemergence at 0.4 oz per acre plus label rates of 2,4-D or MCPA (amine or ester) and surfactant. Volunteer corn: Apply to emerged volunteer corn up to 18" in height. For best results, make FINESSE herbicide application at 0.5 oz per acre preplant or prior to winter wheat emergence. After wheat has emerged, applications are limited to 0.4 oz per acre. Wild buckwheat: For best results, apply FINESSE herbicide preemergence at 0.4 oz per acre to wild buckwheat in the fall or early spring. For postemergence applications, tank mix FINESSE herbicide at 0.4 oz per acre with 2,4-D, MCPA, or dicamba products registered for this use (including Banvel herbicide/Clarity herbicide) and surfactant. Apply after the majority of seedlings have emerged and are actively growing.	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
	0.3	0.0117	0.0023
	0.4	0.0156	0.0031

Note: In certain situations, 0.3 oz of FINESSE herbicide may provide acceptable control of Wild buckwheat. Consult local FMC directions for additional information. Wild radish: For best results, apply FINESSE herbicide at 0.3 to 0.4 oz per acre postemergence.	0.5	0.0195	0.0039
---	-----	--------	--------

TANK MIXTURES

FINESSE herbicide may be tank mixed with other registered herbicides, fungicides, insecticides, or liquid fertilizer. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. If the instructions on the tank mix partner label conflict with this FINESSE herbicide label, DO NOT use in a tank mixture with FINESSE herbicide.

Since tank-mix partners can interfere with FINESSE herbicide dispersion in the spray solution, it is advised that FINESSE herbicide be slurried in a separate container before adding it to the tank mix. FINESSE herbicide must be in suspension in the spray tank before adding companion products.

With 2,4-D (amine or ester) or MCPA (amine or ester)

FINESSE herbicide can be used as a tank-mix treatment with 2,4-D or MCPA herbicides (ester formulations provide best results) after weeds have emerged. For best results, use 0.2 to 0.4 oz of FINESSE herbicide per acre; add 2,4-D or MCPA herbicides to the tank at label rates. Surfactant may be added to the mixture at 0.5 to 1 qt per 100 gal of spray solution; however, adding surfactant may increase the potential for crop injury. DO NOT add a surfactant when FINESSE herbicide plus 2,4-D or MCPA is applied with liquid fertilizer. Apply FINESSE herbicide plus MCPA after the 3 to 5-leaf stage but before boot stage. Apply FINESSE herbicide plus 2,4-D after tillering but before boot stage (refer to the appropriate 2,4-D manufacturer's label). Applying a tank mixture of FINESSE herbicide, 2,4-D, or MCPA and liquid fertilizer when temperatures are below freezing or when the crop is stressed from cold weather just prior to winter dormancy can result in foliar burn and/or crop injury.	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
	0.2 to 0.4	0.0078 to 0.0156	0.0016 to 0.0031

With Dicamba

FINESSE herbicide may be tank mixed with products containing the active ingredient dicamba (including Banvel herbicide) at label rates. Use higher rates when weed infestation is heavy. Nonionic surfactant may be added to the mixture at 0.5 to 1 quart per 100 gal of spray solution (0.125 to 0.25% v/v); however, adding nonionic surfactant may increase the potential for crop injury.

With Diuron

In areas where annual bluegrass, annual ryegrass, corn growwell, green foxtail (pigeongrass) and wild buckwheat are the main weed problems, apply Karmex DF herbicide at label rates plus 0.3 to 0.4 oz per acre FINESSE herbicide preemergence. For best results between 1/2" and 1" of rainfall is needed within 1 to 2 weeks after application. Follow all restrictions and use instructions on the diuron labels. For summer fallow (CO, KS, NE, NM, OK, SD, TX, WY), apply Karmex DF herbicide or Direx 4L herbicide at label rates (both products contain the active ingredient diuron) to wheat stubble or fallow in a tank mix with FINESSE herbicide at 0.2 to 0.3 ounce per acre. Add a Crop Oil Concentrate (COC) at 1 to 2 % v/v or a non-ionic surfactant (NIS) at 0.25 to 0.5 % v/v. Glyphosate products plus AMS may also be added as needed. When using glyphosate products that contain a built-in adjuvant system, add a NIS at 0.25% v/v. Allow at least 90 days after application before planting winter wheat. Refer to the tank mix partners for rates and use instructions.	FINESSE herbicide	Active Ingredient Equivalent	
	Rate (Oz/A)	Chlorsulfuron (Lb. ai/A)	Metsulfuron-methyl (Lb. ai/A)
	0.2 0.3 0.4	0.0078 0.0117 0.0156	0.0016 0.0023 0.0031

With fluroxypyr

FINESSE herbicide may be tank mixed with fluroxypyr containing herbicides for improved control of Kochia (2-4" tall) and other broadleaf weeds at label rates. 2,4-D and MCPA herbicides may be tank mixed with FINESSE herbicide plus fluroxypyr.

With Other Broadleaf Control Products

For improved control of broadleaf weeds, FINESSE herbicide can be tank mixed with other herbicides registered on cereals including Widematch herbicide, Aim® EC herbicide, Huskie® herbicide or Talinor™ herbicide.

With Grass Control Products

For improved control of grass weeds, FINESSE herbicide can be tank mixed with other grass control herbicides registered on cereals including Axial® XL herbicide, Discover® NG herbicide, Everest® 3.0 herbicide or PowerFlex® herbicide.

Weed control antagonism generally does not occur when tank mixing herbicides. However, FMC advises that you first consult your state experiment station, university, or extension agent, Agricultural dealer, or FMC representative as to the potential for antagonism before using the mixture. If no information is available, limit the initial use of FINESSE herbicide and the grass product to a small area.

With Insecticides

FINESSE herbicide may be tank mixed with insecticides registered for use on wheat, barley, and fallow. However, under certain conditions (drought or cold stress while crop is in the 2- to 4-leaf stage), tank mixtures or sequential treatments of FINESSE herbicide and organophosphate insecticides may produce temporary crop yellowing or, in severe cases, crop injury. The potential for crop injury is greatest when there are wide fluctuations in day/night temperatures just prior to or soon after treatment. Read and follow directions on companion product labels and limit first use to a small area. If no symptoms of crop injury appear, larger acreage can be treated.

Restrictions

DO NOT apply FINESSE herbicide within 60 days of crop emergence where an organophosphate insecticide has been applied as an in-furrow treatment, as crop injury may result.

DO NOT use FINESSE herbicide plus malathion, as crop injury may result.

In the Pacific Northwest, DO NOT use FINESSE herbicide with products containing the active ingredient chlorpyrifos, as crop injury may result.

With Fungicides

FINESSE herbicide may be tank mixed with other fungicides whenever the proper timing for herbicide and fungicide treatments coincide.

With Liquid Nitrogen Fertilizer Solution

Liquid nitrogen fertilizer solutions may be used as a carrier in place of water. Run a tank mix compatibility test before mixing FINESSE herbicide in fertilizer solution. If 2,4-D or MCPA is included with FINESSE herbicide and fertilizer mixture, ester formulations tend to be more compatible (See manufacturer's label).

Restrictions

DO NOT add surfactant when using FINESSE herbicide in tank mix with 2,4-D ester or MCPA ester and liquid nitrogen fertilizer solutions.

DO NOT use with liquid fertilizer solutions with a pH less than 3.0.

DO NOT use low rates of liquid fertilizer solution as a substitute for surfactant.

If using low rates of liquid nitrogen fertilizer in the spray solution (less than 50% of the spray solution volume), the addition of surfactant is necessary. When using high rates of liquid nitrogen fertilizer in the spray solution, adding surfactant increases the risk of crop injury. Consult local specifications for details on surfactant addition.

GRAZING

There are no grazing restrictions on FINESSE herbicide.

CROP ROTATION

Before using FINESSE herbicide, carefully consider your crop rotation plans and options. For rotational flexibility, DO NOT treat all of your wheat, barley, or fallow acres at the same time.

MINIMUM ROTATION INTERVALS

Minimum rotation intervals* are determined by the rate of breakdown of FINESSE herbicide applied. FINESSE herbicide breakdown in the soil is affected by soil pH, soil temperature and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase FINESSE herbicide breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow FINESSE herbicide breakdown.

Of these three factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture need to be monitored regularly when considering rotating to other crops.

* The minimum rotation interval represents the period of time from the last FINESSE herbicide application to the anticipated date of the next planting.

SOIL pH LIMITATIONS

FINESSE herbicide may not be used on fields having a soil pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond those specified in the rotation table, and under certain conditions, could injure wheat or barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of FINESSE herbicide. FINESSE herbicide may not be used on soils with a pH below 5.0, as additional crop stress from low pH and aluminum toxicity may result in crop injury.

Checking Soil pH

Before using FINESSE herbicide, determine the soil pH of the field. To obtain a representative pH value, take several samples from different areas of the field between 0" and 4" deep and analyze them separately. Consult local extension publications for additional information on advised soil sampling procedures.

BIOASSAY

A field bioassay must be completed before rotating to any crop not listed (See the Rotation Intervals table), or if the soil pH is not in the specified range, or if the use rate applied is not specified in the table, or if the minimum cumulative precipitation has not occurred since application.

To conduct a field bioassay, grow test strips of the crop or crops you plan to grow the following year in fields previously treated with FINESSE herbicide. Crop response to the bioassay will indicate whether or not to rotate to the crop(s) grown in the test strips.

If a field bioassay is planned, check with your local state agricultural extension service for information detailing the field bioassay procedure.

CEREAL CROPS—ROTATION INTERVALS

Location	Soil pH*	Application Rate (oz/A)	Active Ingredient	Application Rate (Lb. ai/A)	Minimum Rotation Interval (Months)		
					Wheat/Rye/Triticale**	Oat	Barley
AL, AR, DE, GA, IA, IL, IN, KS, KY, LA, MD, MO, MS, NC, NE, NJ, NM, OH, OK, PA, SC, TN, TX, VA	7.9 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	0	10	10
			Metsulfuron-methyl	0.0016 to 0.0031			
	7.9 or lower	0.5	Chlorsulfuron	0.0195	4	10	16
			Metsulfuron-methyl	0.0039			
CO, NE (Panhandle), Southeastern WY	7.9 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	0	10	10
			Metsulfuron-methyl	0.0016 to 0.0031			
ID, OR, WA, MT, ND, SD, and WY (except Southeastern WY)	6.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	0	10	10
			Metsulfuron-methyl	0.0016 to 0.0031			
	6.6 to 7.9	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	0	10	16
			Metsulfuron-methyl	0.0016 to 0.0031			

* See the **Maximum Use Rates** and **Soil pH Limitations** sections of this label.

** For Durum wheat and Wampur variety of Spring Wheat, follow the rotation intervals listed under Barley

NON CEREAL CROPS—ROTATION INTERVALS—NON IRRIGATED LAND

Location		Crop	Soil pH	Application Rate (oz/A)	Active Ingredient	Application Rate (Lb. ai/A)	Cumulative Precipitation (Inches)	Rotation Interval (Months)
State	County or Area							
Colorado	E. of Continental Divide	Field corn, Millets	7.4 or lower 7.5 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	20	11
					Metsulfuron-methyl	0.0016 to 0.0031	45	36
		Grain sorghum	7.5 or lower 7.6 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	45	36
					Metsulfuron-methyl	0.0016 to 0.0031	60	48
		BOLT® technology soybeans**	7.9 or lower***	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	‡	4
					Metsulfuron-methyl	0.0016 to 0.0039		
		STS® & Sulfonylurea Ready soybeans**	7.5 or lower***	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	‡	4†
					Metsulfuron-methyl	0.0016 to 0.0031		
		Grain sorghum	7.2 or lower 7.3 - 7.5***	0.2 to 0.3 0.2 to 0.3	Chlorsulfuron	0.0078 to 0.0117	‡ ‡	4† 8†
					Metsulfuron-methyl	0.0016 to 0.0023		
Idaho*	Northern (Benewah, Bonner, Boundary, Clearwater, Idaho, Koontenai, Latah, Lewis, and Nez Perce counties)	Pea (dry)	6.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	35	24
					Metsulfuron-methyl	0.0016 to 0.0031		
		Lentils	6.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	50	36
					Metsulfuron-methyl	0.0016 to 0.0031		

NON CEREAL CROPS—ROTATION INTERVALS—NON IRRIGATED LAND (CONTINUED)

Location		Crop	Soil pH	Application Rate (oz/A)	Active Ingredient	Application Rate (Lb. ai/A)	Cumulative Precipitation (Inches)	Rotation Interval (Months)
State	County or Area							
Kansas	All areas	Field Corn, Millets	7.4 or lower 7.5 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	20	11
					Metsulfuron-methyl	0.0016 to 0.0031	45	36
		BOLT® technology soybeans**	7.9 or lower***	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	‡	4
					Metsulfuron-methyl	0.0016 to 0.0039		
		STS® & Sulfonylurea Ready soybeans**	7.5 or lower***	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	‡	4†
					Metsulfuron-methyl	0.0016 to 0.0031		
	Central (Generally E. of Highway 183, W. of the Flint Hills)	Grain sorghum Soybeans	7.9 or lower	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	25	14
					Metsulfuron-methyl	0.0016 to 0.0039		
	W. Central and Western (generally W. of Highway 183 to the western edge of Grant, Kearny, Logan, Rawlins, Stevens, Thomas, and Wichita counties)	Grain sorghum	7.5 or lower 7.6 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	21	14
					Metsulfuron-methyl	0.0016 to 0.0031	42	26
		Soybeans	7.5 or lower 7.6 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	40	24
					Metsulfuron-methyl	0.0016 to 0.0031	60	36
	Far Western (In the last tier of counties along the KS/CO border: Cheyenne, Greeley, Hamilton, Morton, Sherman, Stanton, and Wallace)	Grain sorghum Soybeans	7.5 or lower 7.6 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	36	26
					Metsulfuron-methyl	0.0016 to 0.0031	60	48
Western (W. of hwy 183)	Grain sorghum	7.2 or lower 7.3 - 7.5***	0.2 to 0.3 0.2 to 0.3	Chlorsulfuron	0.0078 to 0.0117	‡	4†	
				Metsulfuron-methyl	0.0016 to 0.0023	‡	6†	
Eastern (E. of hwy 183)	Grain sorghum	7.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	‡	4†	
				Metsulfuron-methyl	0.0016 to 0.0031			
Nebraska	All areas	Field Corn, Millets	7.4 or lower 7.5 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	20	11
					Metsulfuron-methyl	0.0016 to 0.0031	45	36
		BOLT® technology soybeans**	7.9 or lower***	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	‡	4
					Metsulfuron-methyl	0.0016 to 0.0039		
		STS® & Sulfonylurea Ready soybeans**	7.5 or lower***	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	‡	4†
					Metsulfuron-methyl	0.0016 to 0.0031		
	S. Central (Franklin, Nuckolls, Thayer, and Webster counties)	Grain sorghum Soybeans	7.9 or lower	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	25	14
					Metsulfuron-methyl	0.0016 to 0.0039		
	Western counties (Chase, Dundy, Frontier, Furnas, Gosper, Harlan, Hayes, Hitchcock, Perkins, Phelps, and Red Willow)	Grain sorghum, Soybeans	7.5 or lower 7.6 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	40	24
					Metsulfuron-methyl	0.0016 to 0.0031	60	36
	Panhandle (Deuel, Garden, and Sheridan counties and all counties W. to the WY border)	Grain sorghum	7.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	45	24
					Metsulfuron-methyl	0.0016 to 0.0031		
	Western (W. of hwy 183)	Grain sorghum	7.2 or lower 7.3 - 7.5***	0.2 to 0.3 0.2 to 0.3	Chlorsulfuron	0.0078 to 0.0117	‡	4†
					Metsulfuron-methyl	0.0016 to 0.0023	‡	6†
Eastern (E. of hwy 183)	Grain sorghum	7.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	‡	4†	
				Metsulfuron-methyl	0.0016 to 0.0031			

NON CEREAL CROPS—ROTATION INTERVALS—NON IRRIGATED LAND (CONTINUED)

Location		Crop	Soil pH	Application Rate (oz/A)	Active Ingredient	Application Rate (Lb. ai/A)	Cumulative Precipitation (Inches)	Rotation Interval (Months)
State	County or Area							
Oklahoma	All areas	Field Corn, Millets	7.4 or lower 7.5 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	20	11
					Metsulfuron-methyl	0.0016 to 0.0031	45	36
		BOLT® technology soybeans**	7.9 or lower***	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	‡	4
					Metsulfuron-methyl	0.0016 to 0.0039		
		STS® & Sulfonylurea Ready soybeans**	7.5 or lower***	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	‡	4†
					Metsulfuron-methyl	0.0016 to 0.0031		
	East of Panhandle	Grain sorghum, Cotton, Mung beans, Soybeans	7.9 or lower	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	25	14
					Metsulfuron-methyl	0.0016 to 0.0039		
	Panhandle	Grain sorghum	7.2 or lower	0.2 to 0.3	Chlorsulfuron	0.0078 to 0.0117	‡	4†
			7.3 - 7.5***		Metsulfuron-methyl	0.0016 to 0.0023	‡	6†
			up to 7.9	up to 0.4	Chlorsulfuron	0.0156	30	25
			Metsulfuron-methyl	0.0031				
All areas except Panhandle	Grain sorghum	7.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	‡	4†	
				Metsulfuron-methyl	0.0016 to 0.0031			
Oregon*	Northeastern counties (Baker, Umatilla, Union, Wallowa)	Pea (dry)	6.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	35	24
					Metsulfuron-methyl	0.0016 to 0.0031		
		Lentils	6.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	50	36
					Metsulfuron-methyl	0.0016 to 0.0031		
	West of the Cascades	Ryegrass (annual and perennial) Crimson Clover	6.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	20	9
					Metsulfuron-methyl	0.0016 to 0.0031		
		Red Clover Snap Beans	6.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	40	15
					Metsulfuron-methyl	0.0016 to 0.0031		
		Field Corn	6.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	60	22
Metsulfuron-methyl	0.0016 to 0.0031							

NON CEREAL CROPS—ROTATION INTERVALS—NON IRRIGATED LAND (CONTINUED)

Location		Crop	Soil pH	Application Rate (oz/A)	Active Ingredient	Application Rate (Lb. ai/A)	Cumulative Precipitation (Inches)	Rotation Interval (Months)	
State	County or Area								
Texas	All areas	Field Corn, Millets	7.4 or lower 7.5 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	20	11	
					Metsulfuron-methyl	0.0016 to 0.0031	45	36	
		BOLT® technology soybeans**	7.9 or lower***	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	‡	4	
					Metsulfuron-methyl	0.0016 to 0.0039			
		STS® & Sulfonylurea Ready soybeans**	7.5 or lower***	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	‡	4†	
					Metsulfuron-methyl	0.0016 to 0.0031			
	Eastern counties (see below)	Grain sorghum, Cotton, Mung beans, Soybeans	7.9 or lower	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	25	14	
					Metsulfuron-methyl	0.0016 to 0.0039			
	The Eastern counties are: Archer, Bell, Bosque, Bowie, Camp, Cass, Clay, Colin, Cooke, Coryell, Dallas, Delta, Denton, Ellis, Falls, Fannin, Franklin, Grayson, Hill, Hood, Hopkins, Hunt, Jack, Johnson, Kaufman, Lamar, Limestone, McLennan, Milam, Montague, Morris, Navarro, Palo Pinto, Parker, Rains, Red River, Robertson, Rockwall, Somervell, Tarrant, Titus, Upshur, Van Zandt, Wichita, Williamson, Wise, Wood, Young								
	Central counties (see below)	Cotton, Grain sorghum	7.9 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	25	14	
					Metsulfuron-methyl	0.0016 to 0.0031			
			7.9 or lower	0.5	Chlorsulfuron	0.0195	46	26	
Metsulfuron-methyl					0.0039				
The Central counties are: Baylor, Callahan, Eastland, Foard, Hardeman, Haskell, Knox, Shackelford, Stephens, Throckmorton, Wilbarger									
Panhandle	Grain sorghum	7.2 or lower 7.3 - 7.5***	0.2 to 0.3 0.2 to 0.3	Chlorsulfuron	0.0078 to 0.0117	‡	4†		
				Metsulfuron-methyl	0.0016 to 0.0023				
		up to 7.9	up to 0.4	Chlorsulfuron	0.0156	30	25		
				Metsulfuron-methyl	0.0031				
All areas except Panhandle	Grain sorghum	7.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	‡	4†		
				Metsulfuron-methyl	0.0016 to 0.0031				
Washington*	Eastern (Asotin, Columbia, Garfield, Pend Oreille, Spokane, Stevens, Walla Walla, Whitman)	Pea (dry)	6.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	35	24	
					Metsulfuron-methyl	0.0016 to 0.0031			
	Lentils	6.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	50	36		
				Metsulfuron-methyl	0.0016 to 0.0031				
Wyoming	Southeastern counties (Platte, Goshen, and Laramie)	Field corn, Millets	7.4 or lower 7.5 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	20	11	
					Metsulfuron-methyl	0.0016 to 0.0031	45	36	
		Grain sorghum	7.5 or lower 7.6 to 7.9	0.2 to 0.4 0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	45	36	
					Metsulfuron-methyl	0.0016 to 0.0031			60
<p>Note: DO NOT plant sorghum grown for hybrid seed production.</p> <p>* In Idaho, Oregon & Washington for peas and lentils, a field bioassay is required if soil pH is above 6.5</p> <p>** BOLT® Technology, Sulfonylurea Ready (SR) & STS® soybeans are varieties that have a high degree of crop non-sensitivity to ALS inhibiting and/or sulfonylurea herbicides. Consult seed provider for confirmation. Under certain conditions (such as drought, prolonged cold weather, pH variability in the fields) temporary discoloration and/or crop injury may occur to above listed soybeans with sulfonylurea non-sensitive traits planted after FINESSE herbicide applications.</p> <p>***Where a CATASTROPHIC CROP LOSS has occurred after a FINESSE herbicide application due to a natural disaster (such as freezing weather, hail damage, insect damage, disease damage), grain sorghum can be planted at 4 months where the soil pH is 7.3 to 7.5 or STS® & Sulfonylurea Ready soybeans can be planted at 4 months where the soil pH is 7.5 to 7.9. These crops will have some level of temporary discoloration and/or crop injury if planted at this reduced interval after FINESSE herbicide application. This potential damage and yield loss is accepted by the grower due to the critical need to get a crop planted after this emergency. Growers not willing to accept this level of potential early season crop injury and yield loss need to follow the standard rotational guidelines in the table above. In some cases, this injury may be severe and may affect the crop growth, development, and yield. The severity of the injury increases with higher pH levels, higher applied FINESSE herbicide rate, drier soil conditions after FINESSE herbicide application and prior to planting the rotational crop, and the shorter the rotational interval.</p> <p>†These intervals may also be used for irrigated land. These intervals DO NOT apply to crops grown for seed.</p> <p>‡Rotation intervals are based on normal precipitation/irrigation amounts. If in a water deficit such as a drought, extend rotation intervals until cumulative rainfall/irrigation reaches the normal range.</p>									

NON CEREAL CROPS—ROTATION INTERVALS—IRRIGATED AND NON IRRIGATED LAND

State	Crop	Soil pH	Application Rate (oz/A)	Active Ingredient	Application Rate (Lb. ai/A)	Rotation Interval* (months)
AL, AR, DE, GA, IL, IN, KY, LA, MD, MS, MO, NC, NJ, OH, PA, SC, TN, VA, WV	BOLT® technology soybeans†	7.9 or lower	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	4
				Metsulfuron-methyl	0.0016 to 0.0039	
	STS®& Sulfonylurea Ready soybeans†	7.9 or lower	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	6
				Metsulfuron-methyl	0.0016 to 0.0039	
	Grain Sorghum, Cotton, Soybeans, Field Corn, Rice	7.9 or lower	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	18
				Metsulfuron-methyl	0.0016 to 0.0039	
	Grain sorghum	7.5 or lower	0.2 to 0.4	Chlorsulfuron	0.0078 to 0.0156	4
				Metsulfuron-methyl	0.0016 to 0.0031	

*Rotation intervals are based on normal precipitation/irrigation amounts. If in a water deficit such as a drought, extend rotation intervals until cumulative rainfall/irrigation reaches the normal range. These intervals DO NOT apply to crops grown for seed.

† BOLT® Technology, Sulfonylurea Ready, & STS® soybeans are varieties that have a high degree of crop non-sensitivity to ALS inhibiting and/or sulfonylurea herbicides. Consult seed provider for confirmation. Under certain conditions (such as drought, prolonged cold weather, pH variability in the fields) temporary discoloration and/or crop injury may occur to above listed soybeans with sulfonylurea non-sensitivity traits planted after FINESSE herbicide applications.

ROTATION INTERVAL TO SOYBEANS WITH THE PLENISH® TRAIT

CROP ROTATION

Minimum Rotation Intervals

Labelled crops may be planted at specified time intervals following application of labelled rates of FINESSE herbicide cereal and fallow herbicide. Soybeans with the PLENISH® trait can be planted 6 months after a labelled application of FINESSE herbicide, where soil pH is 7.9 or lower.

NON CEREAL CROPS -- ROTATION INTERVALS-IRRIGATED AND NON-IRRIGATED LAND

State	Crop	Soil pH	Application Rate (oz/A)	Active Ingredient	Application Rate (Lb. ai/A)	Rotation Interval* (months)
All Registered States	Soybeans with the Plenish® Trait†	7.9 or lower	0.2 to 0.5	Chlorsulfuron	0.0078 to 0.0195	6
				Metsulfuron-methyl	0.0016 to 0.0039	

*Rotational intervals are based on normal precipitation/irrigation amounts. If in a water deficit such as a drought, extend rotation intervals until cumulative rainfall/irrigation reaches the normal range. These intervals do not apply to crops grown for seed.

† Under certain conditions (such as drought, prolonged cold weather, pH variability in fields), temporary discoloration and/or crop injury may occur to soybeans with the Plenish® trait planted after FINESSE herbicide applications.

APPLICATION INFORMATION

PRODUCT MEASUREMENT

FINESSE herbicide is measured using the FINESSE herbicide volumetric measuring cylinder. The degree of accuracy of this cylinder varies by $\pm 7.5\%$. For more precise measurement, use scales calibrated in ounces.

MIXING INSTRUCTIONS

1. Fill the tank 1/4 to 1/3 full of water (If using liquid nitrogen fertilizer solution in place of water, see Tank Mixtures sections for additional details).
2. While agitating, add the required amount of FINESSE herbicide.
3. Continue agitation until the FINESSE herbicide is fully dispersed, at least 5 minutes.
4. Once the FINESSE herbicide is fully dispersed, maintain agitation and continue filling tank with water. FINESSE needs to be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add tank mix partners (if desired) then add the necessary volume of nonionic surfactant. Always add surfactant last.
6. If the mixture is not continuously agitated, settling will occur. If settling occurs, thoroughly re-agitate before using.
7. Apply FINESSE herbicide spray mixture within 24 hours of mixing to avoid product degradation.
8. If FINESSE herbicide and a tank mix partner are to be applied in multiple loads, pre-slurry the FINESSE herbicide in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the

FINESSE herbicide.

DO NOT use FINESSE herbicide with spray additives that reduce the pH of the spray solution to below 3.0.

APPLICATION METHOD

Ground Application

To obtain optimum spray distribution and thorough coverage, use flat-fan or low-volume flood nozzles.

When using flat-fan nozzles, use a spray volume of at least 3 GPA. When using flood nozzles on 30" spacings, use at least 10 GPA, flood nozzles no larger than TK10 (or the equivalent), and a pressure of at least 30 psi. For 40" nozzle spacings, use at least 13 GPA; for 60" spacings, use at least 20 GPA. It is essential to overlap the nozzles 100% for all spacings.

With Raindrop® RA nozzles, DO NOT use less than 20 GPA and overlap nozzles 100%.

Use screens that are 50-mesh or larger.

Aerial Application

Use nozzle types and arrangements that provide optimum spray distribution and maximum coverage at 1 to 5 GPA. Use at least 3 GPA in Idaho, Oregon and Washington.

When applying FINESSE herbicide by air in areas near sensitive crops, use solid-stream nozzles oriented straight back.

Chemigation

DO NOT apply FINESSE herbicide through any type of irrigation system.

SPRAY EQUIPMENT

For specific application equipment, refer to the manufacturer's specifications for additional information on GPA, pressure, speed, nozzle types and arrangements, nozzle heights above the target canopy.

Be sure to calibrate air or ground equipment properly before application. Select a spray volume and delivery system that will ensure thorough coverage and a uniform spray pattern with minimum drift. Use higher spray volumes to obtain better coverage when crop canopy is dense. Avoid swath overlapping, and shut off spray booms while starting, turning, slowing, or stopping, to avoid injury to the crop.

DO NOT make applications using equipment and/or spray volumes or under weather conditions that might cause spray to drift onto nontarget sites. Continuous agitation is required to keep FINESSE herbicide in suspension.

Before Spraying FINESSE herbicide

Spray equipment must be cleaned before FINESSE herbicide is sprayed. Follow the cleanup procedures specified on the labels of previously applied products. If no directions are provided, follow the 6 steps outlined below.

At the End of the Day

When multiple loads of FINESSE herbicide are applied, it is advised that at the end of each day of spraying, the interior of the tank be rinsed with fresh water and then partially filled, and the boom and hoses be flushed. This will prevent the buildup of dried pesticide deposits from accumulating in the application equipment.

After Spraying FINESSE herbicide and before Spraying Crops Other Than Wheat Barley, Triticale, or Fallow

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of FINESSE herbicide as follows:

1. Drain tank; thoroughly rinse spray tanks, boom, and hoses with clean water. Loosen and physically remove any visible deposits.
2. Fill the tank with clean water and 1 gal of household ammonia* (contains at least 3% active ingredient) for every 100 gal of water. Flush the hoses, boom, and nozzles with the cleaning solution. Then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 min. Flush the hoses, boom, and nozzles again with the cleaning solution, and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing ammonia* and water.
4. Repeat step 2.
5. Rinse the tank, boom, and hoses with clean water.
6. If only ammonia is used as a cleaner, the rinsate solution may be applied back to the crop(s) specified on this label. DO NOT exceed the maximum labeled use rate. If other cleaners are used, consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

* Equivalent amounts of an alternate-strength ammonia solution or a cleaner, which dissolves and removes sulfonyleurea herbicide residues can be used in the cleanout procedure. Carefully read and follow the individual cleaner instructions.

Notes:

1. **Caution:** DO NOT use chlorine bleach with ammonia, as dangerous gases will form. DO NOT clean equipment in an enclosed area.

2. Steam-cleaning aerial spray tanks is advised prior to performing the above cleanout procedure to facilitate the removal of any caked deposits.
3. When FINESSE herbicide is tank mixed with other pesticides, all required cleanout procedures need to be examined and the most rigorous procedure needs to be followed.
4. In addition to this cleanout procedure, all preapplication cleanout guidelines on subsequently applied products need to be followed as per the individual labels.
5. Where routine spraying practices include shared equipment frequently being switched between applications of FINESSE herbicide and applications of other pesticides to FINESSE herbicide-sensitive crops during the same spray season, it is advised that a sprayer be dedicated to FINESSE herbicide to further reduce the chance of crop injury.

MANDATORY SPRAY DRIFT MANAGEMENT

Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 3 feet above the ground or crop canopy.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Aerial Applications:

- Do not release spray at a height greater than 10 feet above the vegetative canopy, unless a greater application height is necessary for pilot safety.
- For applications prior to the emergence of crops and target weeds, applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- For all other applications, applicators are required to use a Medium or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 65% of the wingspan for airplanes or 75% of the rotor blade diameter for helicopters.
- Applicators must use one-half swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

Boom-less Ground Applications:

- Applicators are required to use a Medium or coarser droplet size (ASABE S572.1) for all applications.
- Do not apply when wind speeds exceed 10 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT MANAGEMENT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to

reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom should remain level with the crop and have minimal bounce.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift. When applying aurally to crops, do not release spray at a height greater than 10 ft above the crop canopy, unless a greater application height is necessary for pilot safety.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

HANDHELD TECHNOLOGY APPLICATIONS:

- Take precautions to minimize spray drift.

BOOM-LESS GROUND APPLICATIONS:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

DRIFT CONTROL ADDITIVES

Using product compatible drift control additives can reduce drift potential. When a drift control additive is used, read and carefully observe cautionary statements and all other information on the additive's label. If using an additive that increases viscosity, ensure that the nozzles and other application equipment will function properly with a viscous spray solution. Preferred drift control additives have been certified by the Council of Producers & Distributors of Agrotechnology (CPDA).

IDENTIFICATION INFORMATION FOR PRODUCTS REFERENCED IN THIS LABEL

REGISTERED PRODUCTS REFERENCED IN THIS LABEL		
Product Name	Active Ingredient(s)	EPA Registration Number
Aim® EC herbicide	carfentrazone-ethyl	279-3241
Ally® Extra SG herbicide (with Totalsol® soluble granules)	thifensulfuron-methyl, tribenuron-methyl, metsulfuron-methyl	279-9603
Ally® XP herbicide	metsulfuron-methyl	279-9575
Amber® Custom-Pak™ herbicide	traisulfuron	100-768
Axial® XL herbicide	pinoxaden	100-1256
Banvel® herbicide	dicamba	66330-276
Clarity® herbicide	dicamba	7969-137
Direx® 4L herbicide	Diuron	66222-54
Discover® NG herbicide	clodinafop-propargyl	100-1173
Everest® 3.0 herbicide	flucarbazone-sodium	66330-429
Glean® XP herbicide	chlorsulfuron	279-9600
Huskie® herbicide	pyrasulfotole, bromoxynil	264-1023
Karmex® DF herbicide	diuron	66222-51
PowerFlex® herbicide	pyroxsulam	62719-569
Talinor™ herbicide	bicyclopyrone, bromoxynil	100-1570
Widematch herbicide	clopyralid, fluroxypyr	62719-512

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store product in original container only. Do not contaminate water, other pesticides, fertilizer, food or feed in storage. Store in a cool, dry place.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by disposal. Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.

Nonrefillable Plastic and Metal Containers (Capacity Equal to or Less Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers (Capacity Greater Than 50 Pounds): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Plastic and Metal Containers, e.g., Intermediate Bulk Containers [IBC] (Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down): Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Nonrefillable Paper or Plastic Bags, Fiber Sacks including Flexible Intermediate Bulk Containers (FIBC) or Fiber Drums with Liners: Nonrefillable container. Do not reuse or refill this container. Completely empty paper or plastic bag, fiber sack or drum liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer for recycling if available or dispose of empty paper or plastic bag, fiber sack or fiber drum and liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

Refillable Fiber Drums with Liners: Refillable container (fiber drum only). Refilling Fiber Drum: Refill this fiber drum with FINESSE herbicide containing chlorsulfuron and metsulfuron methyl only. Do not reuse this fiber drum for any other purpose. Cleaning before refilling is the responsibility of the refiller.

Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Disposing of Fiber Drum and/or Liner: Do not reuse this fiber drum for any other purpose other than refilling (see preceding). Cleaning the container (liner and/or fiber drum) before final disposal is the responsibility of the person disposing of the container. Offer the liner for recycling if available or dispose of liner in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner. To clean the fiber drum before final disposal, completely empty the fiber drum by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application or manufacturing equipment. Then offer the fiber drum for recycling if available or dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

All Other Refillable Containers: Refillable container. Refilling Container: Refill this container with FINESSE herbicide containing chlorsulfuron and metsulfuron methyl only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use the container, contact FMC at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container, contact FMC at the number below for instructions. Disposing of Container: Do not reuse this container for any other purpose other than refilling (see preceding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Outer Foil Pouches of Water Soluble Packets (WSP): Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken. If the outer pouch contacts the formulated product in any way, the pouch must be triple rinsed with clean water. Add the rinsate to the spray tank and dispose of the outer pouch as described previously.

Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact CHEMTREC (Transportation and Spills) at 1-800-424-9300, day or night.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

FMC, Ally, Aim, Finesse, Glean and TotalSol are trademarks of FMC Corporation or an affiliate.

Huskie is a trademark of Bayer CropScience

Raindrop is a trademark of GP Companies

Amber, Axial, Discover, and Talinor are trademarks of Syngenta Group Company

PowerFlex, & Widematch are trademarks of the Dow AgroSciences LLC

Banvel and Clarity are trademarks of BASF Corporation

Everest is a trademark of Arysta Lifesciences North America LLC

Karmex and Direx are trademarks of Adama Group Company

SL-4009 071219 07-15-19

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

Notice: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and, to the extent consistent with applicable law, Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been used contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or FMC, and, to the extent permitted by applicable law, Buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Condition of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.