



**Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170.**

**Not for sale, distribution, or use in Nassau or Suffolk Counties in New York State.**

<b>ACTIVE INGREDIENTS:</b>	
2,4-D, dimethylamine salt	18.79%
Mecoprop-p, dimethylamine salt	6.80%
Dicamba, dimethylamine salt	3.02%
Sulfentrazone	0.67%
<b>INERT INGREDIENTS:</b>	70.72%
	<b>TOTAL 100.00%</b>

**THIS PRODUCT CONTAINS:**  
 1.40 lb. 2,4-dichlorophenoxyacetic acid equivalent per gallon or 15.66%.  
 0.50 lb. (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per gallon or 5.62%.  
 0.22 lb. 3,6-dichloro-o-anisic acid equivalent per gallon or 2.52%.  
 0.06 lb. N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl]methanesulfonamide per gallon or 0.67%.  
 Isomer Specific by AOAC Methods.

## KEEP OUT OF REACH OF CHILDREN DANGER – PELIGRO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find some one to explain it to you in detail.)



**READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.**

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

**DANGER:** Corrosive. Causes irreversible eye damage. Do not get in eyes, on skin or on clothing. Harmful if absorbed through skin or if swallowed.

#### Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are natural rubber, natural rubber blends and laminates. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, and other handlers must wear\*:

- protective eyewear,
- long-sleeved shirt and long pants,
- shoes and socks, plus
- chemical-resistant gloves, and
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

\*Applicators may choose not to wear protective eyewear when dilution is with water only and dilution rates are greater (higher) than 5:1 or greater (higher) than 5 parts of water to 1 part of product.

See engineering controls for additional requirements.

#### User Safety Requirements

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Control Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### First Aid

<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or on clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

#### Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls worn over short-sleeve shirt and short pants,
- chemical-resistant footwear plus socks
- chemical-resistant gloves made of any waterproof material,
- chemical-resistant headgear for overhead exposure, and
- protective eyewear.

#### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Reentry Statement:** Do not enter or allow people (or pets) to enter the treated area until sprays have dried.



**Droplet Size**

Use only Medium or coarser spray nozzles according to ASAE (S 572) definition of standard nozzles or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

**Wind Speed**

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind.

**Temperature Inversions**

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

**Susceptible Plants**

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

**Other State and Local Requirements**

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

**Equipment**

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

**Application Schedules**

Apply this product to broadleaf weeds that are young and actively growing for the best results. Spring and fall treatments under adequate soil moisture conditions are preferred to the summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective. Fall applications provide improved control for emerged winter annuals and perennials such as henbit, chickweed, clover and ground ivy.

Sequential broadcast applications or follow-up applications as spot treatments at a 30-day interval are suggested for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment, or any time broadleaf weeds are actively growing.

Extremes in environmental conditions e.g. temperature and moisture, soil conditions, and cultural practices may affect the activity of this product. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicide symptoms is delayed, and weeds hardened off by drought are less susceptible to this product.

**For newly seeded areas:**

Delay application of this product to grass seedlings until after the second mowing.

**For newly sodded, sprigged, or plugged areas:**

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

**Reseeding interval:**

Treated areas may be reseeded 3 weeks after application.

**Irrigation:**

Do not apply this product through any type of irrigation system. Rainfast in as little as 6 hours. Do not apply this product immediately before rainfall or irrigation. If possible, do not irrigate or water the turfgrass within 6 to 24 hours after application. If dry conditions exist, a scheduled irrigation or watering 24 hours before and 24 hours after application is suggested.

**Mowing:**

Delay mowing 2 days before and until 2 days after the application of this product.

**How Much To Use**

**Use Rates and Spray Volumes:** Generally, the lower application rates within the specified range of Table 1 will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

**BROADCAST TREATMENT:**

Table 1. Use rates for sod farms, ornamental lawns and turfgrass.		
Species	Rate	Spray Volume
<b>Cool-season Turf</b>		
Kentucky bluegrass, annual bluegrass, annual ryegrass, perennial ryegrass, tall fescue, red or fine leaf fescues, creeping bentgrass and colonial bentgrass (excluding golf greens and tees). Mixtures of cool-season species in non-cropland areas established for roadside vegetation management or for low maintenance. (Kentucky bluegrass, tall fescue, smooth brome grass & orchardgrass)	3.25 to 4 Pints/Acre (1.2 to 1.5 fl. oz./ 1,000 sq. ft.)	10 to 220 Gallons/Acre (0.23 to 5.0 Gallons/ 1,000 sq. ft.)
<b>Warm-season Turf</b>		
Hybrid Bermudagrass, common Bermudagrass, centipedegrass*, zoysiagrass, bahiagrass and buffalograss.	2.75 to 3.25 Pints/Acre (1.0 to 1.2 fl. oz./ 1,000 sq. ft.)	10 to 220 Gallons/Acre (0.23 to 5.0 Gallons/ 1,000 sq. ft.)
*For centipedegrass use lower rates within specified range until turfgrass tolerance to injury can be determined.		
Do not apply this product to warm-season turfgrass during spring green-up or in the fall during the transition period between active growth and dormancy. <b>Dormant turf:</b> This product may be applied to fully dormant bermudagrass, fully dormant centipedegrass, fully dormant zoysiagrass, and fully dormant bahiagrass. <b>New York:</b> Only one application per year of this product is allowed. This product is not allowed to be sold, distributed or used in Nassau or Suffolk Counties.		

**SPOT TREATMENT:**

**WITH HAND OPERATED SPRAYERS (INCLUDING BACKPACK SPRAYERS, COMPRESSION SPRAYERS, AND KNAPSACK SPRAYERS):**

- Apply any time the emerged broadleaf weeds are actively growing.
- Calibration and proper application are essential when using this product.
- Uniform applications are essential when using this product. Over application or rates above those specified on this label including excessive overlaps of this product can cause turf injury.
- Hand-held techniques: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Spray wands fitted with flat fan tips should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motions result in uneven coverage.
- Follow-up applications as spot treatments at a 30-day interval are suggested for more mature weeds, for dense infestations, and for adverse environmental conditions.
- **For cool-season turfgrass listed in Table 1:** Mix 1.2 to 1.5 fl. oz. of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq. ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.
- **For warm-season turfgrass listed in Table 1:** Mix 1.0 to 1.2 fl. oz. of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq. ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible. For centipedegrass use lower rates within specified range until turfgrass tolerance to injury can be determined.

**Limitations on spot treatments for ornamental turfgrass:**

Spot treatment is defined as a treatment area no greater than 1,000 sq. ft. per acre. The maximum application rate is 1.5 fl. oz. per 1,000 sq. ft. per application (0.25 lb. MCP-P acid equivalent per acre). The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

**TOUGH WEED CONTROL:**

**IN INDUSTRIAL OR LOW MAINTENANCE AREAS:**

In addition to weeds listed in Table 4, Surge Broadleaf Herbicide For Turf can be used for certain tough-to-control weeds. Mixed stands of Kentucky bluegrass, tall fescue, smooth brome grass, orchardgrass and reed canarygrass may be treated. Applications to non-cropland areas (roadsides and rights-of-way) are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Treatments of Surge Broadleaf Herbicide For Turf may injure or kill legumes including clovers (sweet, yellow, red, crimson, alsike, hop, white), lespedezas, trefoils and vetches.

Biennial and perennial weeds may require follow-up or sequential treatments.

**Table 2. Rates of product per acre for specific hard-to-control weeds with ground equipment.**

Weed Types	Troublesome weeds such as:	Amount of product, pints/acre <sup>1</sup>	Spray Volume	When to Apply
Annual Broadleaf	ivyleaf morningglory, redroot pigweed, cocklebur, sunflower, velvetleaf (butterprint)	2.5 to 4.0 pints/acre	10 to 220 gal./Acre (0.23 to 5.0 gal. per 1,000 sq. ft.)	Spring or fall during active growth.
Biennial	Bull thistle, musk thistle, common burdock	4.0 to 5.0 pints/acre	10 to 220 gal./Acre (0.23 to 5.0 gal. per 1,000 sq. ft.)	Spring or fall during seedling to rosette stage.
Perennial	Burclover, Canada thistle, field bindweeds, English daisy, hoary cress (whiteweed), Veronica (corn speedwell), wild violet	4.0 to 5.0 pints/acre	10 to 220 gal./Acre (0.23 to 5.0 gal. per 1,000 sq. ft.)	Spring or fall during bud to bloom stage.

**Footnote 1:** Use the lower rate within the range specified for highly favorable plant growing conditions and when broadleaf weeds are less than 6 inches in height. Use the higher rate within the range specified for tall vegetation, dense canopies, weeds beyond the suggested growth stage, or during adverse conditions.

**New York:** Only one application per year of this product is allowed. This product is not allowed to be sold, distributed or used in Nassau or Suffolk Counties.

**LIMITATIONS:**

**Table 3. Limitations on broadcast applications to ornamental turfgrass, sod farms, roadsides and rights-of-way (noncropland)**

Use Site	Maximum Rate per Application	Maximum Number of Applications per Year	Minimum Interval Between Applications	Maximum Seasonal Rate
Ornamental turfgrass	4.0 pints/A	2	30 Days	8.0 pints/A
Sod farms	4.0 pints/A	2	21 Days	8.0 pints/A
Non-cropland: roadsides and rights-of-way	5.0 pints/A	2	30 Days	10.0 pints/A

For ornamental turfgrass only, do not exceed the maximum seasonal rate of 8.0 pints per acre per season, excluding spot treatments. For non-cropland: Do not exceed the maximum seasonal rate of 10 pints per acre per season.

**Spray volumes with ground equipment:**

- For Surge Broadleaf Herbicide For Turf alone, a spray volume of 10.0 gallons per acre (gpa) is suggested.
- For tank mixtures of Surge Broadleaf Herbicide For Turf, spray volume of 20.0 gallons per acre is suggested or use the specified spray volume of the companion product(s).

**Broadleaf Weeds Controlled**

Surge Broadleaf Herbicide For Turf will control or suppress the following broadleaf weeds. Apply any time the emerged broadleaf weeds are susceptible.

**Table 4. Broadleaf Weeds**

Aster, white heath & white prairie	Dogfennel	Knotweed
Bedstraw	Dollarweed (*pennywort)	Lambsquarters
Beggarweed, creeping	False dandelion	Lawn burweed
Bindweed	(*spotted catsear & common catsear)	Lespedeza, common
Black medic	Field bindweed	Mallow, common
Broadleaf plantain	(*morningglory & creeping jenny)	Matchweed
Buckhorn plantain	Field oxeye-daisy	Mouseear chickweed
Bull thistle	(*creeping oxeye)	Nutsedge** (yellow)
Burdock, common	Filaree, whitestem & redstem	Old world diamond flower
Buttercup, creeping	Florida betony	Oxalis (*yellow woodsorrel & creeping woodsorrel)
Carpetweed	Florida pusley	Parsley-piert
Chickweed, common	Ground ivy	Pennsylvania smartweed
Chicory	Groundsel	Pepperweed
Cinquefoil	Hawkweed	Pigweed
Clover	Healall	Pineappleweed
Curly dock	Henbit	Plantain
Dandelion	Innocence (Blue-eyed Mary)	
Dayflower		
Deadnettle		
Dock		

(cont. on next column)

**Table 4. Broadleaf Weeds (cont.)**

Poison ivy	Shepherdspurse	Wild lettuce
Poison oak	Spurge	Wild mustard
Prickly lettuce	Thistle	Wild onion
(*compass plant)	Virginia buttonweed	Wild strawberry
Puncturevine	White clover (*Dutch clover, honeysuckle clover, white trefoil, & purplewort)	Yarrow
Purple cudweed	Wild carrot	Yellow rocket
Purslane	Wild garlic	
Ragweed	Wild geranium	
Redweed		
Red sorrel (*sheep sorrel)		

\*Synonyms

\*\*Suppression only when nutsedge is young and actively growing.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**For Plastic Containers - Nonrefillable with capacities equal to or less than 5 gallons:**

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

OR

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For Plastic Containers - Nonrefillable with capacities greater than 5 gallons:**

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

OR

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**For Refillable Containers:**

**CONTAINER HANDLING:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Container cleaning: Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

